

## THE MEDICAL CENTER AT BOWLING GREEN

# Cancer Program

ANNUAL OUTCOMES REPORT 2019

### QUALITY MEASURES

2016 Outcome Analysis – Cancer Program Practice Profile Reports (CP<sup>3</sup>R)

Site	Criteria	Measure	Med Center Health %	Commission on Cancer Benchmark
COLON	At least 12 regional lymph nodes are removed and pathologically examined for resected colon cancer (Quality Improvement)	12RLN	97.40%	85%
	Adjuvant chemotherapy is recommended, or administered within 4 months (120 days) of diagnosis for patients under the age of 80 with AJCC stage III (lymph node positive) colon cancer (Accountability)	ACT	91.70%	Not Defined
RECTUM	Preoperative chemo and radiation are administered for clinical AJCC T3N0, T4N0, or Stage III; or Postoperative chemo and radiation are administered within 180 days of diagnosis for clinical AJCC T1-2N0 with pathologic AJCC T3N0, T4N0, or Stage III; or treatment is recommended; for patients under the age of 80 receiving resection for rectal cancer (Quality Improvement)	RECRCT	100%	85%

#### Resources:

The Medical Center at Bowling Green Cancer Registry Statistical Data  
National Cancer Database (NCDB)



Med Center Health.

The Medical Center at Bowling Green