

the medical center at bowling green

ANNUAL OUTCOMES REPORT 2018

Med Center Health The Medical Center at Bowling Green

QUALITY MEASURES

Site	Criteria	Measure	Ratio	%	on Cancer Benchmark
BREAST	Radiation is administered within 1 year (365 days) of diagnosis for women under the age of 70 receiving breast conservation surgery for breast cancer (Accountability)	BCSRT	23/23	100%	90%
	Tamoxifen or third generation aromatase inhibitor is recommended or administered within 1 year (365 days) of diagnosis for women with AJCC T1c or stage IB-III hormone receptor positive breast cancer (Accountability)	ΗT	40/41	97.60%	90%
	Radiation therapy is recommended or administered following any mastectomy within 1 year (365 days) of diagnosis of breast cancer for women with >= 4 positive regional lymph nodes (Accountability)	MASTRT	5/7	71.40%	90%
	Image or palpation-guided needle biopsy to the primary site is performed to establish diagnosis of breast cancer (Quality Improvement)	nBx	93/106	87.70%	80%
	Breast conservation surgery rate for women with AJCC clinical stage 0, I, or II breast cancer (Surveillance)	BCS	30/66	45.50%	Not Defined
	Combination chemotherapy is recommended or administered within 4 months (120 days) of diagnosis for women under 70 with AJCC T1cN0, or stage IB - III hormone receptor negative breast cancer (Accountability)	MAC	3/3	100%	Not Defined
	Systemic chemotherapy is administered within 4 months to day preoperatively or day of surgery to 6 months postoperatively, or it is recommended for surgically resected cases with pathologic lymph node-positive (pN1) and (pN2) NSCLC (Quality Improvement)	LCT	11/11	100%	85%
	Surgery is not the first course of treatment for cN2, M0 lung cases (Quality Improvement)	LNoSurg	20/22	90.90%	85%
	At least 10 regional lymph nodes are removed and pathologically examined for AJCC Stage IA, IB, IIA, and IIB resected NSCLC (Surveillance)	10RLN	28/53	52.80%	Not Defined



Commission